

(19) World Intellectual Property
Organization
International Bureau



(43) International Publication Date
29 April 2004 (29.04.2004)

PCT

(10) International Publication Number
WO 2004/035723 A1

- (51) International Patent Classification⁷: C11D 9/26, 9/24, 10/04, 3/18, 3/20
- (21) International Application Number: PCT/GB2003/004422
- (22) International Filing Date: 13 October 2003 (13.10.2003)
- (25) Filing Language: English
- (26) Publication Language: English
- (30) Priority Data: 0223847.5 12 October 2002 (12.10.2002) GB
- (71) Applicant (*for all designated States except US*): RECKITT BENCKISER INC [US/US]; Morris Corporate Center IV, 399 Interpace Parkway, Parippany, NJ 07054 (US).
- (72) Inventors; and
- (75) Inventors/Applicants (*for US only*): CHEUNG, Tak, Wal [GB/US]; Reckitt Benckiser Inc, One Phillips Parkway, Montvale, NJ 07645 (US). HARIHARAN, Ramakrishnan [US/US]; Reckitt Benckiser Inc, One Phillips Parkway, Montvale, NJ 07645 (US). LU, Robert, Zhong [US/US]; Reckitt Benckiser Inc, One Phillips Parkway, Montvale, NJ 07645 (US). NGUYEN, Vu [VN/US]; Reckitt Benckiser Inc, One Phillips Parkway, Montvale, NJ 07645 (US). CHRISTMAS, Delford [US/US]; Reckitt Benckiser Inc, One Phillips Parkway, Montvale, NJ 07645 (US).
- (74) Agents: MCKNIGHT, John, Crawford et al.; Reckitt Benckiser plc, Group Patents Department, Dansom Lane, Hull, Yorkshire HU8 7DS (GB).
- (81) Designated States (*national*): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BR, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, HR, HU, ID, IL, IN, IS, JP, KE, KG, KP, KR, KZ, LC, LK, LR, LS, LT, LU, LV, MA, MD, MG, MK, MN, MW, MX, MZ, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RU, SC, SD, SE, SG, SK, SL, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, YU, ZA, ZM, ZW.
- (84) Designated States (*regional*): ARIPO patent (GH, GM, KE, LS, MW, MZ, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IT, LU, MC, NL, PT, RO, SE, SI, SK, TR), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).
- Declarations under Rule 4.17:**
- *as to applicant's entitlement to apply for and be granted a patent (Rule 4.17(ii)) for all designations*
 - *of inventorship (Rule 4.17(iv)) for US only*
- Published:**
- *with international search report*
- For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.*

(54) Title: DISINFECTANT CLEANING COMPOSITIONS

(57) Abstract: The present invention provides a non-cationic antimicrobial agent containing composition which blooms when added to water. The compositions have good cleaning, disinfecting and bloom properties.

WO 2004/035723 A1

JC06 Rec'd PCT/PTO 05 APR 2005

5

DISINFECTANT CLEANING COMPOSITIONS

The present invention is directed to concentrated cleaning and/or disinfecting compositions which bloom when diluted in water.

Cleaning compositions are commercially important products and enjoy a wide field of utility in assisting in the removal of dirt and grime from surfaces, especially those characterized as useful with hard surfaces. One particular category of cleaning compositions are those which provide a blooming effect. Such an effect may be described as the change of the water's appearance from essentially colorless and transparent to that of a milky white or milky yellowish white, cloudy appearance upon the addition of an amount of the cleaning composition. This effect is also sometimes referred to as the "break". Such blooming is a highly desirable in such pine oil type cleaning compositions as consumer/end user expectations associate cleaning effectiveness/disinfection with the extent and degree of this blooming upon formation of a cleaning composition. Such an effect is particularly known and generally associated with pine oil type cleaning compositions which typically include one or more of the following identifying characteristics: containing an amount of one or more resins or oils derived from coniferous species of trees; containing natural fragrances or synthetic fragrance compositions which are intended to mimic the scent of one or more resins or oils derived from coniferous species of trees; a color ranging from colorless to a deep amber, deep amber yellow or deep amber reddish color; generation of a milky or cloudy appearance when diluted with water in dilutions useful for cleaning applications. Such pine oil type cleaning compositions are generally provided in a concentrated composition which is subsequently diluted with water by an end user/consumer to form a cleaning composition therefrom.

Thus, the present invention provides a hard surface cleaning concentrate composition comprising:

a) from about 0.05 to about 15 wt%, preferably about 0.1 to about 5 wt%, and more preferably from about 0.5 to about 5 wt% of a non-cationic antimicrobial agent;

b) from about 0.1 to about 20 wt% and more preferably from about 0.5 to about 10 wt% of a water soluble organic solvent;

5 c) from about 0.1 to about 30 wt%, preferably from about 1 to about 20 wt%, and more preferably from about 2 to about 15 wt% of an anionic soap surfactant;

d) from about 0.1 to about 15 wt%, preferably from about 0.1 to about 10 wt%, and more preferably from about 0.5 to about 5 wt% of an hydrocarbon diluent;

10 e) from about 0.001 to about 10 wt%, preferably from about 0.1 to about 5 wt%, and more preferably from about 0.5 to about 5 wt% of pine oil which is at least 60% terpene alcohols;

f) optionally, from about 0 to about 10 wt% of optional materials selected from dyes, colorants, pH stabilizers and buffers, non-ionic surfactants, fragrance/fragrance enhancers which do not interfere with the pine oil constituent, viscosity modifiers, insect repellants, and light stabilizers; and

15 g) the balance being water.

In some preferred embodiments, one or more insect repellants is present.

Preferably, the non-cationic antimicrobial agent is selected from pyrrithiones, dimethyldimethylol hydantoin, methylchloroisothiazolinone/methylisothiazolinone sodium sulfite, sodium bisulfite, imidazolidinyl urea, diazolidinyl urea, benzyl alcohol, 2-bromo-2-nitropropane-1,3-diol, formalin (formaldehyde), iodopropenyl butylcarbamate, chloroacetamide, methanamine, methyldibromonitrile glutaronitrile, glutaraldehyde, 5-bromo-5-nitro-1,3-dioxane, phenethyl alcohol, o-phenylphenol/sodium o-phenylphenol, sodium hydroxymethylglycinate, polymethoxy bicyclic oxazolidine, dimethoxane, thimersal

25 dichlorobenzyl alcohol, captan, chlorphenenesin, dichlorophene, chlorbutanol, glyceryl laurate, halogenated diphenyl ethers, phenolic compounds, mono- and poly-alkyl and aromatic halophenols, resorcinol and its derivatives, bisphenolic compounds, benzoic esters (parabens), halogenated carbanilides, 3-trifluoromethyl-4,4'-dichlorocarbanilide, and 3,3',4-trichlorocarbanilide. More preferably, the non-cationic antimicrobial agent is a mono- and

30 poly-alkyl and aromatic halophenol selected from the group p-chlorophenol, methyl p-

chlorophenol, ethyl p-chlorophenol, n-propyl p-chlorophenol, n-butyl p-chlorophenol, n-amyl p-chlorophenol, sec-amyl p-chlorophenol, n-hexyl p-chlorophenol, cyclohexyl p-chlorophenol, n-heptyl p-chlorophenol, n-octyl p-chlorophenol, o-chlorophenol, methyl o-chlorophenol, ethyl o-chlorophenol, n-propyl o-chlorophenol, n-butyl o-chlorophenol, n-amyl o-chlorophenol, tert-amyl o-chlorophenol, n-hexyl o-chlorophenol, n-heptyl o-chlorophenol, o-benzyl p-chlorophenol, o-benzyl-m-methyl p-chlorophenol, o-benzyl-m, m-dimethyl p-chlorophenol, o-phenylethyl p-chlorophenol, o-phenylethyl-m-methyl p-chlorophenol, 3-methyl p-chlorophenol, 3,5-dimethyl p-chlorophenol, 6-ethyl-3-methyl p-chlorophenol, 6-n-propyl-3-methyl p-chlorophenol, 6-iso-propyl-3-methyl p-chlorophenol, 2-ethyl-3,5-dimethyl p-chlorophenol, 6-sec-butyl-3-methyl p-chlorophenol, 2-iso-propyl-3,5-dimethyl p-chlorophenol, 6-diethylmethyl-3-methyl p-chlorophenol, 6-iso-propyl-2-ethyl-3-methyl p-chlorophenol, 2-sec-amyl-3,5-dimethyl p-chlorophenol 2-diethylmethyl-3,5-dimethyl p-chlorophenol, 6-sec-octyl-3-methyl p-chlorophenol, p-chloro-m-cresol, p-bromophenol, methyl p-bromophenol, ethyl p-bromophenol, n-propyl p-bromophenol, n-butyl p-bromophenol, n-amyl p-bromophenol, sec-amyl p-bromophenol, n-hexyl p-bromophenol, cyclohexyl p-bromophenol, o-bromophenol, tert-amyl o-bromophenol, n-hexyl o-bromophenol, n-propyl-m,m-dimethyl o-bromophenol, 2-phenyl phenol, 4-chloro-2-methyl phenol, 4-chloro-3-methyl phenol, 4-chloro-3,5-dimethyl phenol, 2,4-dichloro-3,5-dimethylphenol, 3,4,5,6-terabromo-2-methylphenol, 5-methyl-2-pentylphenol, 4-isopropyl-3-methylphenol, para-chloro-meta-xilenol, dichloro meta xylenol, chlorothymol, and 5-chloro-2-hydroxydiphenylmethane.

Preferably, the water soluble organic solvents are short chain alcohols.

Preferably, for the anionic soap surfactants, alkyl metal soaps and alkylaryl sulfonates are preferred with alkyl metal soaps, such as sodium or potassium castor oil soap, being preferred.

Preferably, the hydrocarbon diluent is an aromatic hydrocarbon diluent, preferably selected from Shellsolv AB, Aromatic 150, Aromatic 200 (naphthalene depleted), Aromatic 200, Aromatic 100, and HAN 857.

According to certain especially preferred embodiments, the concentrate compositions may be characterized in that when the concentrate compositions are diluted at a ratio of 1 part

to 100 parts water at 20°C. or 40°C. the resultant mixture exhibits a light transmittance loss of at least 50%. In particularly preferred embodiments the concentrate compositions do not form a gel at usual storage conditions (room temperature, approx. 20°C.), and exhibit a satisfactory blooming effect when added to a larger volume of water where such water is
5 either at room temperature or at a higher temperature, particularly at approx. 40°C.

According to preferred embodiments, the concentrate compositions cause a drop in transmitted light through water of at least 20%, more desirably at least about 30% and more when used to form a cleaning composition therefrom, particularly at a dilution of 1 part cleaning concentrate to 100 parts water relative to the transmittance of water, which is
10 established to be 100%. According to particularly preferred embodiments the concentrate compositions cause a drop in transmitted light through water of at least about 40% when added to water at 40°C., and also cause a drop in transmitted light through water of at least 40%, more desirably at least 50%, when added to water at 20°C. Most preferred are compositions which exhibit a drop in transmitted light of at least 70% when added to water at
15 20°C., and which exhibit a drop in transmitted light of at least 60% when added to water at 40°C. wherein the dilutions of concentrate composition to water is 1:100.

The antimicrobial agents of the present invention are non-cationic in order to avoid interaction with the anionic soap surfactant of the invention. Given below are examples of non-cationic antimicrobial agents which are useful in the present invention: pyrithiones
20 (especially zinc pyrithione which is also known as ZPT), dimethyldimethylol hydantoin (Glydant®), methylchloroisothiazolinone/methylisothiazolinone (Kathon CG®), sodium sulfite, sodium bisulfite, imidazolidinyl urea (Germall 115®), diazolidinyl urea (Germaill II®), benzyl alcohol, 2-bromo-2-nitropropane-1,3-diol (Bronopol®), formalin (formaldehyde), iodopropenyl butylcarbamate (Polyphase P100®), chloroacetamide,
25 methanamine, methyldibromonitrile glutaronitrile (1,2-Dibromo-2,4-dicyanobutane or Tektamer®), glutaraldehyde, 5-bromo-5-nitro- 1,3-dioxane (Bronidox®), phenethyl alcohol, o-phenylphenol/sodium o-phenylphenol, sodium hydroxymethylglycinate (Suttocide A®), polymethoxy bicyclic oxazolidine (Nuosept C®), dimethoxane, thimersal dichlorobenzyl alcohol, captan, chlorphenenesin, dichlorophene, chlorbutanol, glyceryl laurate, halogenated
30 diphenyl ethers like 2,4,4'-trichloro-2'-hydroxy-diphenyl ether (Triclosan® or TCS), 2,2'-

dihydroxy-5,5'-dibromo-diphenyl ether, phenolic compounds like phenol, 2-methyl phenol, 3-methyl phenol, 4-methyl phenol, 4-ethyl phenol, 2,4-dimethyl phenol, 2,5-dimethyl phenol, 3,4-dimethyl phenol, 2,6-dimethyl phenol, 4-n-propyl phenol, 4-n-butyl phenol, 4-n-amyl phenol, 4-tert-amyl phenol, 4-n-hexyl phenol, 4-n-heptyl phenol, mono- and poly-alkyl and

5 aromatic halophenols such as p-chlorophenol, methyl p-chlorophenol, ethyl p-chlorophenol, n-propyl p-chlorophenol, n-butyl p-chlorophenol, n-amyl p-chlorophenol, sec-amyl p-chlorophenol, n-hexyl p-chlorophenol, cyclohexyl p-chlorophenol, n-heptyl p-chlorophenol, n-octyl p-chlorophenol, o-chlorophenol, methyl o-chlorophenol, ethyl o-chlorophenol, n-propyl o-chlorophenol, n-butyl o-chlorophenol, n-amyl o-chlorophenol, tert-amyl o-

10 chlorophenol, n-hexyl o-chlorophenol, n-heptyl o-chlorophenol, o-benzyl p-chlorophenol, o-benzyl-m-methyl p-chlorophenol, o-benzyl-m, m-dimethyl p-chlorophenol, o-phenylethyl p-chlorophenol, o-phenylethyl-m-methyl p-chlorophenol, 3-methyl p-chlorophenol, 3,5-dimethyl p-chlorophenol, 6-ethyl-3-methyl p-chlorophenol, 6-n-propyl-3-methyl p-chlorophenol, 6-iso-propyl-3-methyl p-chlorophenol, 2-ethyl-3,5-dimethyl p-chlorophenol,

15 6-sec-butyl-3-methyl p-chlorophenol, 2-iso-propyl-3,5-dimethyl p-chlorophenol, 6-diethylmethyl-3-methyl p-chlorophenol, 6-iso-propyl-2-ethyl-3-methyl p-chlorophenol, 2-sec-amyl-3,5-dimethyl p-chlorophenol 2-diethylmethyl-3,5-dimethyl p-chlorophenol, 6-sec-octyl-3-methyl p-chlorophenol, p-chloro-m-cresol, p-bromophenol, methyl p-bromophenol, ethyl p-bromophenol, n-propyl p-bromophenol, n-butyl p-bromophenol, n-amyl p-

20 bromophenol, sec-amyl p-bromophenol, n-hexyl p-bromophenol, cyclohexyl p-bromophenol, o-bromophenol, tert-amyl o-bromophenol, n-hexyl o-bromophenol, n-propyl-m,m-dimethyl o-bromophenol, 2-phenyl phenol, 4-chloro-2-methyl phenol, 4-chloro-3-methyl phenol, 4-chloro-3,5-dimethyl phenol, 2,4-dichloro-3,5-dimethylphenol, 3,4,5,6-terabromo-2-methylphenol, 5-methyl-2-pentylphenol, 4-isopropyl-3-methylphenol, para-chloro-meta-

25 xlenol, dichloro meta xlenol, chlorothymol, 5-chloro-2-hydroxydiphenylmethane, resorcinol and its derivatives including methyl resorcinol, ethyl resorcinol, n-propyl resorcinol, n-butyl resorcinol, n-amyl resorcinol, n-hexyl resorcinol, n-heptyl resorcinol, n-octyl resorcinol, n-nonyl resorcinol, phenyl resorcinol, benzyl resorcinol, phenylethyl resorcinol, phenylpropyl resorcinol, p-chlorobenzyl resorcinol, 5-chloro 2,4-

30 dihydroxydiphenyl methane, 4'-chloro 2,4-dihydroxydiphenyl methane, 5-bromo 2,4-

dihydroxydiphenyl methane, and 4'-bromo 2,4-dihydroxydiphenyl methane, bisphenolic compounds like 2,2'-methylene bis (4-chlorophenol), 2,2'-methylene bis (3,4,6-trichlorophenol), 2,2'-methylene bis (4-chloro-6-bromophenol), bis (2-hydroxy-3,5-dichlorophenyl) sulphide, and bis (2-hydroxy-5-chlorobenzyl)sulphide, benzoic esters

5 (parabens) like methylparaben, propylparaben, butylparaben, ethylparaben, isopropylparaben, isobutylparaben, benzylparaben, sodium methylparaben, and sodium propylparaben, halogenated carbanilides (e.g., 3,4,4'-trichlorocarbanilides (Triclocarban® or TCC), 3-trifluoromethyl-4,4'-dichlorocarbanilide, 3,3',4-trichlorocarbanilide, etc.). The phenol based non-cationic antimicrobials are preferred. The non-cationic antimicrobial agent
10 in the present invention is present in an amount from about 0.05 to about 15 wt%, preferably from about 0.1 to about 5 wt%, and more preferably from about 0.5 to about 5 wt%.

Water soluble organic solvents include short chain (1 to 4 carbon atoms) alcohols, glycol ethers, and mixtures thereof. Such water soluble organic solvents provide effective solubilization of many types of greases and fats which may be encountered in soils, as well
15 as being useful in the solubilization of the pine oil in water, without substantially interfering with the blooming and scent characteristics of the compositions according to the present invention. Of course two or more organic solvents may be used as the organic solvent constituent according to the invention. Examples of short chain alcohols include ethanol, propanol, and isopropanol. Examples of such useful glycol ethers include propylene glycol
20 methyl ether, dipropylene glycol methyl ether, tripropylene glycol methyl ether, propylene glycol n-propyl ether, diethylene glycol methyl ether, ethylene glycol butyl ether, diethylene glycol methyl ether, and mixtures thereof. The short chain alcohols are preferred. The water soluble organic solvent constituent may be present in the concentrate compositions in amounts of from about 0.1 to about 20 wt%, preferably in amounts of from about 0.5 to about
25 10 wt%.

Anionic soap surfactants represent the primary detergent component in the present compositions of interest. This class of surfactants includes ordinary alkali metal soaps such as the sodium, potassium, ammonium and alkanol-ammonium salts of higher fatty acids containing from about 8 to about 24 carbon atoms and preferably from about 10 to about 20
30 carbon atoms. Suitable fatty acids can be obtained from natural sources such as, for instance,

plant or animal esters (*e.g.*, palm oil, coconut oil, babassu oil, soybean oil, castor oil, tallow, whale and fish oils, grease, lard, and mixtures thereof). The fatty acids also can be synthetically prepared (*e.g.*, by the oxidation of petroleum, or by the Fischer-Tropsch process). Resin acids are suitable such as rosin and those resin acids in tall oil. Naphthenic acids are also suitable. Sodium and potassium soaps can be made by direct saponification of the fats and oils or by the neutralization of the free fatty acids which are prepared in a separate manufacturing process. Particularly useful is the sodium or potassium salt of the mixtures of fatty acids derived from castor oil, *i.e.*, sodium or potassium castor oil soap.

The anionic soap surfactant can also comprise traditional anionic surfactants such as alkali metal salts, ammonium salts, amine salts, aminoalcohol salts or the magnesium salts of one or more of the following compounds: alkyl sulfates, alkyl ether sulfates, alkylamidoether sulfates, alkylaryl polyether sulfates, alkylaryl sulfates, alkylaryl sulfonates, monoglyceride sulfates, alkylsulfonates, alkylamide sulfonates, alkylarylsulfonates, olefinsulfonates, paraffin sulfonates, alkyl sulfosuccinates, alkyl ether sulfosuccinates, alkylamide sulfosuccinates, alkyl sulfosuccinamate, alkyl sulfoacetates, alkyl phosphates, alkyl ether phosphates, acyl sarconsinates, acyl isethionates, and N-acyl taurates. Generally, the alkyl or acyl radical in these various compounds comprise a carbon chain containing 12 to 20 carbon atoms. Of these anionic surfactants, the alkylaryl sulfonates are preferred. From the anionic soap surfactants contemplated herein, the alkyl metal soaps, such as sodium or potassium castor oil soap, are preferred.

The anionic soap surfactant in the present invention comprises an amount from about 0.1 to about 30 wt%, preferably from about 1 to about 20 wt%, and more preferably from about 2 to about 15 wt%.

Another component of the present invention is a hydrocarbon diluent, which is preferably an aromatic hydrocarbon diluent. For example, the hydrocarbon diluent may be Shellsolv AB, Aromatic 150, Chemical Abstract Service (CAS) Registry No. 64742-94-5; Aromatic 200 (naphthalene depleted), CAS Registry No. 64742-94-5; Aromatic 200, Aromatic 100, CAS Registry No. 64742-95-6; and HAN (heavy atmospheric naphtha) 857, CAS Registry No. 64742-06-9. Aromatic 200 is generally preferred. Shellsolv AB is commercially available from Shell Chemical Company. Aromatic 150, Aromatic 100,

Aromatic 200 (naphthalene depleted), Aromatic 200, and HAN 857 are commercially available from ExxonMobil. Other suitable aromatic solvents are available from Koch Chemical, Ashland, Amoco, and Texaco as well as the Isopar, Exxsol and Norpar aliphatic, isoparaffins and paraffins from ExxonMobil. The hydrocarbon diluent is preferably present
5 in the present composition in an amount of about 0.1 to about 15 wt%, preferably in an amount of from about 0.1 to about 10 wt% and more preferably in an amount of from about 0.5 to about 5 wt%.

The compositions according to the present invention also comprise a pine oil constituent. Pine oil is an organic solvent, and is a complex blend of oils, alcohols, acids,
10 esters, aldehydes and other organic compounds. These include terpenes which include a large number of related alcohols or ketones. Some important constituents include terpineol, which is one of three isomeric alcohols having the basic molecular formula $C_{10}H_{17}OH$. One type of pine oil, synthetic pine oil, will generally have a specific gravity, at 15.5°C. of about 0.9300, which is lower than the two other grades of pine oil, namely steam distilled and sulfate pine
15 oils, and will generally contain a higher content of turpentine alcohols. Other important compounds include alpha- and beta-pinene (turpentine), abietic acid (rosin), and other isoprene derivatives.

Particularly useful pine oils which are presently commercially available include Unipine® 60 (from Union Camp, believed to contain approximately 60% terpene alcohols),
20 Unipine® S-70 and Unipine® S-70 (both are believed to contain approximately 70% terpene alcohols), Unipine® S and Unipine® 80 (both are believed to contain approximately 80% terpene alcohols), Unipine® 85 (which is believed to contain approximately 85% terpene alcohols), Unipine® 90 (which is believed to contain approximately 90% terpene alcohols), as well as Alpha Terpineol 90 (which is believed to contain approximately 100% terpene
25 alcohols). Further useful pine oils include Glidco® Pine Oil™ 60 (available from Glidco Organics Corp., Jacksonville, Fla., believed to contain approximately 60% terpene alcohols), Glidco® Pine Oil 140 (believed to contain approximately 70% terpene alcohols), Glidco® Pine Oil 80 (believed to contain approximately 80% terpene alcohols), Glidco® Pine Oil 150 (believed to contain approximately 85% terpene alcohols); Glidco® Terpene SW (believed to
30 contain approximately 75% terpene alcohols); as well as Glidco® Terpineol 350 (believed to

contain approximately 100% terpene alcohols). Other products which can contain up to 100% pure alpha-terpineol, may also be used in the present invention.

Preferred as the pine oil constituent of the present invention are pine oil preparations which comprise at least about 60% terpene alcohols, and more preferably those which
5 comprise at least about 80% terpene alcohols.

The pine oil constituent may be present in the concentrate compositions in amounts of from about 0.001 to about 20 wt%, preferably in amounts of from about 0.1 to about 15 wt% and more preferably from about 0.5 to about 10 wt%.

As the concentrate compositions are aqueous, water forms a major constituent. Water
10 is added in order to provide 100 wt% of the concentrate composition. The water may be tap water, but is preferably distilled and/or deionized water. If the water is tap water, it is preferably appropriately filtered in order to remove any undesirable impurities such as organics or inorganics, especially mineral salts which are present in hard water which may thus interfere with the operation of the other constituents of the invention, as well as any
15 other optional components of the liquid concentrates according to the invention.

Water is added in amounts which are sufficient to form the concentrated compositions which amount is sufficient to ensure the retention of a substantially clear characteristic when produced as a concentrate, but at the same time ensuring good blooming upon the addition of the concentrated composition to a further amount of water, or upon the addition of further
20 water to the concentrate. This amount may be readily determined by first mixing measured amount of the non-water constituents in a suitably sized vessel and then during stirring adding water. Generally, water is present in the concentrate compositions in amounts in excess of about 50 wt%, preferably in amounts of in excess of about 70 wt%, but most preferably in amount of between 70-80 wt% based on the total weight of the concentrate
25 compositions according to the invention.

As noted previously, the concentrate compositions according to the invention may include further optional, but advantageously included constituents.

Useful optional constituents are one or more coloring agents which find use in modifying the appearance of the concentrate compositions and enhance their appearance
30 from the perspective of a consumer or other end user. Known coloring agents, may be


incorporated in the compositions in effective amount to improve or impart to concentrate compositions an appearance characteristic of a pine oil type concentrate composition, such as a color ranging from colorless to yellow or yellow/brown color with or without fluorescent ingredients. Such a coloring agent or coloring agents may be added in any useful amount in a conventional fashion, *i.e.*, admixing to a concentrate composition or blending with other constituents used to form a concentrate composition. However, other colors atypical of pine oil type and/or lemon oil type cleaning concentrates may be used as well. Known art light stabilizer constituents useful in pine oil type compositions may also be added, particularly wherein coloring agents are used in a composition. As is known to the art, such light stabilizers act to retain the appearance characteristics of the concentrate compositions over longer intervals of time.

Further useful optional constituents which may in some cases be desirably included in the inventive compositions include rheology modifying agents such as thickeners.

Other conventional additives known to the art but not expressly enumerated here may also be included in the compositions according to the invention. By way of non-limiting example these may include fragrance/fragrance adjusters (which do not interfere with the pine oil constituent of the present invention (can generally be present in an amount of from about 0.000001 to about 1.5 wt% of the total concentrate, preferably non-pine oil based)), pH adjusters, pH buffering agents, non-ionic surfactants and insect repellants. Such non-ionic surfactants denoted here are conventionally known; examples are described in McCutcheon's Detergents and Emulsifiers, North American Edition, 2001; Kirk-Othmer, Encyclopedia of Chemical Technology, 4th Ed., Vol. 23, pp. 478-541, the contents of which are herein incorporated by reference. Such optional constituents should be selected so to have little or no detrimental effect upon the desirable characteristics of the present invention, namely the blooming behavior, cleaning efficacy, and disinfectant activity, low toxicity as provided by the inventive compositions. Generally the total weight of such further conventional additives may comprise up to 10 wt% of a concentrated composition formulation.

In certain preferred instances, one of the optional constituents, insect repellants, may be a required component of an inventive composition.

Examples of insect repellants which can be used include those such as essential oils (such as, for example, oils of anise, citrus, aniseed, roses, mint, camphor, lemon, orange, rosemary, wintergreen, thyme, lavender, cloves, hops, tea tree, citronella, wheat, barley, lemongrass, cedar leaf, cedarwood, palmarosa, vetiver, sage, lime, juniper berry, cinnamon, fleagrass, geranium, sandalwood, violet, ginger, bergamot, marjoram, pepper, jasmine, chamomile, petitgrain, cranberry, eucalyptus, vervain, peppermint, gum benzoin, basil, fennel, fir, balsam, menthol, ocmea origanum, hydastis carradensis, berberidaceae daceae, ratanhiaie, curcuma longa, *Mentha arvensis* (Cornmint), *Mentha spicata* (American Spearmint), *Mentha cardica* (Scotch Spearmint), (-)-Limonene, (+)-Limonene, (-)-Carvone, Linalool, Alpha and Beta -Terpineol, Fencholic acid, Borneol iso Borneol, Bornyl acetate and iso Bornyl acetate. Also included in this class of natural essential oils are the key chemical components of the plant oils which have been found to provide the antimicrobial benefit. These chemicals include, but are not limited to anethol, catechole, camphene, pinocarvone, cedrol, thymol, eugenol, eucalyptol, ferulic acid, farnesol, hinokitiol, tropolone, limonene, menthol, methyl salicylate, carvacol, terpineol, verbenone, berberine, ratanhiaie extract, caryophellene oxide, citronellic acid, curcumin, nerolidol and geraniol), N,N-diethyl-m-toluamide, diethyl phthalate, dimethyl phthalate, dibutyl phthalate, 2-Hydroxyethyl-n-octyl sulfide, N-Octyl bicycloheptene dicarboximide, Hexahydrodibenzofuran carboxaldehyde, Di-n-propyl isocinchomerate, 2-Ethyl-1,3-hexanediol, 2-(n-butyl)-2-ethyl-1,3-propanediol, Dibutyl succinate, Piperonyl butoxide, Pyrethrum, certain fragrances, those agents listed in, for example, United States Patent No. 6,451,844, n-alkylneoalkanamides (such as those described in U.S. Pat. Nos. 5,006,562; 5,015,665; 5,143,900; 5,182,304; 5,182,305; 5,258,408; 5,391,578; and 5,434,189; the disclosures of which are incorporated herein by reference) where, for example, n-alkylneoalkanamide can have a formula of



wherein R_1 , R_2 , and R_3 are alkyl groups and the sum of the carbon atoms therein is from 6 to 12, and wherein R_4 is either a hydrogen atom or an alkyl group having one or two carbon atoms and wherein R_5 is an alkyl group having one to three carbon atoms.

When present, the insect repellant is present in the formulations of the present invention in an amount of from about 0.1 to about 5 wt%.

What is to be understood by the term "concentrate" and "concentrate composition" in this specification and claims is the pre-consumer dilution and composition of the cleaning composition which is the essentially the form of the product prepared for sale to the consumer or other end user with required product shelf life. Such a consumer or other end user would then normally be expected to dilute the same with water to form a cleaning composition. It is to be understood however that nothing in this invention would bar its use as cleaning composition without any further dilution and it may be used in the concentrations in which it was prepared for sale. Similarly, what is to be understood by the term "cleaning compositions" are the water diluted compositions which are expected to be prepared by the consumer or other end user by mixing a measured amount of the "concentrate" with water in order to form an appropriately diluted cleaning composition which is suitable for use in cleaning applications, especially in the cleaning of hard surfaces.

It is also to be understood, that proportions of one or more constituents have been and generally are referred to as percent by weight or as parts by weight based on a measure of 100% by weight, unless otherwise indicated.

As generally denoted above, the formulations according to the invention include both cleaning compositions and concentrates as outlined above which differ only in the relative proportion of water to that of the other constituents forming such formulations. While the concentrated form of the cleaning compositions find use in their original form, they are more frequently used in the formation of a cleaning composition therefrom. Such may be easily prepared by diluting measured amounts of the concentrate compositions in water by the consumer or other end user in certain weight ratios of concentrate:water, and optionally, agitating the same to ensure even distribution of the concentrate in the water. As noted, the concentrate may be used without dilution, i.e., in concentrate:water concentrations of 1:0, to extremely dilute dilutions such as 1:2000. Desirably, the concentrate is diluted in the range of

1:0.1-1:1000, preferably in the range of 1:10-1:500 but most preferably in the range of 1:50-1:200. The actual dilution selected is in part determinable by the degree and amount of dirt and grime to be removed from a surface(s), the amount of mechanical force imparted to remove the same, as well as the observed efficacy of a particular dilution. Generally better
5 results and faster removal is to be expected at lower relative dilutions of the concentrate in water.

Examples

Preparation of Example Formulations:

Exemplary formulations illustrating certain preferred embodiments of the inventive
10 compositions and described in more detail in Table I below were formulated generally in accordance with the following protocol.

Into a suitably sized vessel, a measured amount of the non-cationic antimicrobial agent (*e.g.*, PCMX, DCMX) was provided after which the remaining organic based components (*e.g.*, hydrocarbon diluent, pine oil, water soluble organic solvent, dye (if any)).
15 These components were stirred until the non-cationic antimicrobial agent was fully dissolved. The anionic soap surfactant is then added to the organic solution and the components are stirred until the mixture is homogeneous. The remaining amount of water is then added to the organic/anionic soap surfactant mixture and the components are then mixed until the solution is clear and homogeneous. Mixing of the various components at the various times
20 can range from 5 minutes to 120 minutes, depending on the volume of the mixture; the amount of time can vary so long as particular solutions appeared to be homogeneous. The exemplary compositions were readily pourable, and retained well mixed characteristics (*i.e.*, stable mixtures) upon standing for extended periods, even in excess of 90 days. The exemplary compositions are also stable at elevated temperatures up to 120°F for several
25 weeks.

Examples of inventive formulations are shown in Table 1 below.

Table 1							
Components	Ex. 1	Ex. 2	Ex. 3	Ex. 4	Ex. 5	Ex. 6	Ex. 7
Aromatic 200	4.00	1.50	1.50	0.00	1.50	1.00	0.30
Pine Oil 80		1.00	1.00	2.00	1.00	2.00	3.50
PCMX ¹	10.00	1.50	1.50	1.50	0.00	1.40	
DCMX ²					1.50		
OBPCP							1.50
IPA ³	4.00	2.00	2.00	3.00	2.00		3.00
Na COS ⁴	12.93	20.00	20.00	35.00	20.00	35.00	20.00
SDS ⁵	0.80						
SLS ⁶						3.00	
NP-9						0.60	
Insect repellent ⁷			0.40	4.00	0.40	5.00	
Dye (1%)		0.25					
Water, DI	68.27	73.75	73.60	54.50	73.60	52.00	71.70
Total	100.01	100.00	100.00	100.00	100.00	100.00	100.00

Table 1						
Components	Ex. 8	Ex. 9	Ex. 10	Ex. 11	Ex. 12	Ex. 13
Aromatic 200	1.50	0.30	0.50	1.50	1.50	0.00
Aromatic 200 (naphthalene depleted)						0.40
Pine Oil 80 or 85	1.00	3.00	2.00	0.80	1.60	2.00
PCMX	1.40	0.00	0.00	1.50	1.50	1.50
OBPCP ⁸		1.50	1.50			
IPA	2.00	3.00	3.00	2.00	2.00	3.00
Na COS	18.00	18.00	20.00	20.00	20.00	35.00
NP-9 ⁹			1.00			
Insect repellent				0.40	0.40	0.40
Fragrance Oils ¹⁰	0.25			0.20	0.40	
Water, DI	76.10	74.20	72.00	73.60	72.60	54.10
Total	100.00	100.00	100.00	100.00	100.00	100.00

¹ p-chloro-m-xylene

5 ² dichloro meta xylenol

³ isopropanol

⁴ sodium castor oil soap (40%)

⁵ sodium dodecyl benzene sulfonate (40%)

⁶ sodium lauryl sulfate (30%)

10 ⁷ dimethyl phthalate

⁸ ortho-benzyl-para-chlorophenol

⁹ nonylphenol ethoxylates (9 moles EO)

¹⁰ lemongrass oil or eucalyptus oil

15 The compositions of the present invention were evaluated for bloom, cleaning and disinfecting properties.

The formulations according to Ex. 1 through Ex. 13 are clear (although some examples contain dye but these dyes do not affect the clarity of the compositions), but when diluted at ratios of 1 part to 100 parts or 200 parts of water at both 20° C. and 35° C., in the as mixed aqueous dilutions are expected to achieve the targeted loss of light transmittance of about 40% and more. Ex. 2, with and without insect repellent, was tested for bloom. In certain regions of the world where cold water is used for cleaning purposes, formulations of the present invention can achieve a loss of light transmission greater than 90%.

Light transmittance values closer to zero indicate improved blooming behavior. The protocol for evaluating light transmittance is described as follows: The results of the light transmittance evaluation was determined as a percentage of light transmitted through a sample of a particular aqueous dilution wherein the transmission of a like sample of water is assigned a percentage of 100%. Testing was performed by preparing a 1:100 or 1:200 dilution of the example formulation:water, (tap water) after which the sample was mixed for 30 seconds and a transmittance reading was taken using a Brinkman model PC801 dipping probe colorimeter, which was set at 620 nm to determine the light transmission of each of the samples. Readings were taken at water temperatures of 20° C. and at 35° C. were evaluated, as well as the reference (pure tap water) sample used to calibrate the colorimeter to the reference 100% light transmission sample outlined above. The results are shown in Table 2 below.

Table 2		
Composition	1:100	1:200
In water at 20°C		
Ex. 2 w/IR ¹	0.5	1.3
Ex. 2 w/o IR	0.4	1.1
In water at 35°C		
Ex. 2 w/IR	34.6	50.6
Ex. 2 w/o IR	31.3	61.7

¹ IR is an insect repellent

These results provide an empirical evaluation of the degree of transparency of a diluted example formulation wherein 0% indicates complete opacity and 100% the transparency of the sample. Accordingly, a lower reported light transmittance value of a particular aqueous dilution provided a more desirable indication of the blooming

characteristic of the particular aqueous dilution. It can be noted that in some instances, warm water will be "soft" and cold water will be "hard." In the above table, the warm (35°C) water was treated (or "soft") and thus the blooming characteristics are somewhat reduced.

However, if non-softened warm water were used, similar blooming characteristics to those

5 seen in cold water would be achieved.

Cleaning Evaluations

Cleaning evaluations were performed in accordance with the testing protocol outlined according to ASTM D4488 A2 Test Method, which evaluated the efficacy of the cleaning

10 compositions on masonite wallboard samples painted with wall paint. The soil applied was a greasy soil sample containing vegetable oil, food shortening and animal fat. The sponge (water dampened) of a Gardner Abrasion Tester apparatus was squirted with a 15 gram sample of a tested cleaning composition, and the apparatus was cycled 10 times. The evaluation of cleaning compositions was "paired" with one side of each of the test samples

15 treated with a composition according to the invention, and the other side of the same sample treated with a comparative example's composition, thus allowing a "side-by-side"

comparison to be made. Each of these tests were duplicated on 5 wallboard tiles and the results statistically analyzed and the averaged results reported on Table 3, below. The

20 cleaning efficacy of the tested compositions was evaluated utilizing a Minolta Chroma Meter CF-110, with Data Processor DP-100, which evaluated spectrophotomic characteristics of the sample. The results are reported on Table 3, following.

Table 3	
Composition	Average % Soil Removal
Ex. 2	90.05
Dettol Multi-Purpose Disinfectant from Reckitt Benckiser	91.45
Domex Disinfectant Cleaner from Lever Johnson	57.64
Domex 2-in-1 Phenolic Cleaner from Lever Johnson	69.55
Lizol Pineol Shakti Disinfectant Cleaner from Reckitt Benckiser	87.25

With respect to the results reported on Table 3 a value of "100" is indicative of total soil removal and a "0" value is indicative no soil removal. As can be seen from the results of Table 3, the cleaning efficacy of the composition according to the invention generally provided superior results or were on parity with those of known art cleaning products.

5

Evaluation of Antimicrobial Efficacy

Representative compositions of the present invention were evaluated in order to evaluate their antimicrobial efficacy against *Staphylococcus aureus* (gram positive type pathogenic bacteria) (ATCC 6538), and *Salmonella choleraesuis* (gram negative type pathogenic bacteria) (ATCC 10708). The testing was performed generally in accordance with the protocols outlined in "Use-Dilution Method" as promulgated by the Association of Official Analytical Chemists (AOAC).

The results of the AOAC Use-Dilution Test Method indicates the number of test substrates wherein the tested organism remains viable after contact for 10 minutes with a test disinfecting composition/total number of tested substrates (plates) evaluated in accordance with the AOAC Use-Dilution Test. Thus, a result of "0/10" indicates that of 10 test substrates bearing the test organism and contacted for 10 minutes in a test disinfecting composition, 0 test substrates had viable (live) test organisms at the conclusion of the test. Such a result is excellent, illustrating the excellent disinfecting efficacy of the tested composition.

Results of the antimicrobial testing are indicated on Table 4, below. The reported results indicate the number of test plates with live test organisms/number of test plates tested for each example formulation and organism tested.

Table 4		
Composition (dilution)	<i>S.aureus</i>	<i>S.choleraesuis</i>
Ex. 3 (1:50)	0/10	Not tested
Ex. 7 (1:20)	0/10	0/10

25

Composition Ex. 3 with added dye was evaluated at three dilutions (1:60; 1:90; and 1:100 compositions:water) using the Biomek® 2000 Laboratory Automation Workstation together with the BioWorks Operating System (available from Beckman Coulter Inc., Fullerton, CA) with the above organisms at a concentration of 9 logs. The Biomek simulates

a microbial reduction suspension test. One part of organism suspension (*Staphylococcus aureus* or *Salmonella choleraesuis*) is added to certain dilutions of Ex. 3 in an appropriate container. Deionized water (DI H₂O) was used a control. The organism and sample are then mixed thoroughly for 15 seconds. Serial tenfold dilutions are carried out in a neutralizing
 5 broth. The diluted samples are then incubated for 24-48 hours at 35-37°C. Thereafter, surviving organisms are quantified and Log10 reduction, as a measurement of organism survivors are calculated as follows:

$$\text{Log10 Reduction} = (\text{Log10 Survivors/DI H}_2\text{O Control}) - (\text{Log10 Survivors/Sample})$$

10 For this test with a contact time of 15 second, a Log10 reduction value of 3 or greater against both organisms is a good indication of acceptable performance (i.e., broad spectrum antimicrobial activity). The results of this evaluation are summarized in Table 5 below.

15

Table 5			
Organism/dilution	1:60	1:90	1:100
<i>S.aureus</i>	>6 log	6.7	3.99
<i>S.choleraesuis</i>	>6 log	4.03	2.7

Various compositions were tested for insect repellency (against American cockroach (*Periplaneta americana*) on either vinyl flooring or mosaic flooring. In either using vinyl
 20 flooring or mosaic flooring, a 100 cm x 100 cm square was treated with the compositions at two concentrations: neat and at 5 ml composition/1000 ml water. The compositions were applied to the surface and allowed to dry for 30 minutes.

The treated surface (either vinyl or mosaic) was placed in one half of a suitably sized tray with a similar sized untreated surface placed in the other half of the tray. At the center
 25 of each surface (treated and untreated) were placed two 30 cm x 30 cm sheets of Formica® to act as a cockroach harborage. The Formica® sheet in the treated section was treated at the same rate as the treated surface.

Fifty cockroaches (15 adults, 35 nymphs) were placed in the middle of the tray. No food or water was supplied to the cockroaches.

At the end of thirty minutes, 3 hours, 6 hours, and 12 hours, the number of cockroaches under each Formica® harborage was counted.

Each assay was repeated three times with three replicates each time. The data are shown below in Tables 6, 6A, 7, and 7A.

5

Table 6 - Formulations tested Neat on Vinyl Flooring									
Ex.	The number of cockroaches present in each harborage of treated and untreated vinyl flooring surfaces in different time intervals (each value is the mean of three individual observations)								
	Trial No.	Time in hours							
		30 minutes		3 hours		6 hours		12 hours	
		T	Un. T	T	Un. T	T	Un. T	T	Un. T
Ex. 5	I	4	14	8	20	14	20	26	24
	II	6	16	15	22	16	25	16	32
	III	3	20	12	23	16	24	18	29
Ex. 4	I	3	19	7	23	15	26	26	24
	II	1	16	10	24	18	22	24	26
	III	1	18	13	21	19	24	20	29
Ex. 3	I	8	19	12	22	18	23	27	18
	II	4	17	17	21	16	27	20	26
	III	6	16	14	22	17	21	22	24

Table 6A - Formulations tested 5 ml Ex./1000 ml water on Vinyl Flooring									
Ex.	The number of cockroaches present in each harborage of treated and untreated vinyl flooring surfaces in different time intervals (each value is the mean of three individual observations)								
	Trial No.	Time in hours							
		30 minutes		3 hours		6 hours		12 hours	
		T	Un. T	T	Un. T	T	Un. T	T	Un. T
Ex. 5	I	8	12	22	15	25	22	31	18
	II	10	15	22	18	22	24	23	24
	III	9	16	18	21	25	23	20	26
Ex. 4	I	11	15	17	22	23	20	31	19
	II	10	13	19	26	20	28	20	28
	III	10	20	12	26	18	26	20	28
Ex. 3	I	13	15	19	22	20	22	24	24
	II	15	16	26	22	25	23	20	27
	III	12	18	23	18	27	19	20	20

Table 7 - Formulations tested Neat on Mosaic Flooring									
Ex.	The number of cockroaches present in each harborage of treated and untreated mosaic flooring surfaces in different time intervals (each value is the mean of three individual observations)								
	Trial No.	Time in hours							
		30 minutes		3 hours		6 hours		12 hours	
		T	Un. T	T	Un. T	T	Un. T	T	Un. T
Ex. 5	I	2	18	8	24	14	26	23	27
	II	1	14	9	26	17	28	24	26
	III	0	16	11	28	20	24	21	29
Ex. 4	I	6	12	10	19	15	21	25	25
	II	5	18	16	23	18	24	22	28
	III	2	21	11	24	17	23	24	25
Ex. 3	I	7	18	14	23	16	23	21	29
	II	6	14	17	27	17	27	19	26
	III	4	16	13	21	18	21	24	23

Table 7A - Formulations tested 5 ml Ex./1000 ml water on Mosaic Flooring									
Ex.	The number of cockroaches present in each harborage of treated and untreated mosaic flooring surfaces in different time intervals (each value is the mean of three individual observations)								
	Trial No.	Time in hours							
		30 minutes		3 hours		6 hours		12 hours	
		T	Un. T	T	Un. T	T	Un. T	T	Un. T
Ex. 5	I	12	15	18	22	19	22	22	24
	II	10	16	19	26	22	26	24	26
	III	9	21	21	26	23	26	20	26
Ex. 4	I	9	14	20	19	20	22	20	24
	II	10	19	22	24	25	24	25	24
	III	12	21	21	21	21	25	22	25
Ex. 3	I	11	18	24	19	24	21	24	23
	II	16	16	27	22	25	23	22	24
	III	14	20	23	22	23	22	20	23

We claim:

1. A hard surface cleaning concentrate composition comprising:
 - a) from about 0.05 to about 10 wt% of a non-cationic antimicrobial agent;
 - 5 b) from about 1 to about 20 wt% of a water soluble organic solvent;
 - c) from about 1 to about 20 wt% of an anionic soap surfactant;
 - d) from about 1 to about 15 wt% of a hydrocarbon diluent;
 - e) from about 0.001 to about 20 wt% of pine oil which is at least 60% terpene alcohols;
 - 10 f) optionally, from about 0 to about 10 wt% of optional materials selected from dyes, colorants, pH stabilizers and buffers, non-ionic surfactants, fragrance/fragrance enhancers, viscosity modifiers, insect repellants, and light stabilizers; and
 - g) the balance being water.
- 15 2. The cleaning concentrate of claim 1 wherein the non-cationic antimicrobial agent is selected from pyrrithiones, dimethyldimethylol hydantoin, methylchloroisothiazolinone/methylisothiazolinone sodium sulfite, sodium bisulfite, imidazolidinyl urea, diazolidinyl urea, benzyl alcohol, 2-bromo-2-nitropropane-1,3-diol, formalin (formaldehyde), iodopropenyl butylcarbamate, chloroacetamide,
20 methanamine, methyldibromonitrile glutaronitrile, glutaraldehyde, 5-bromo-5-nitro-1,3-dioxane, phenethyl alcohol, o-phenylphenol/sodium o-phenylphenol, sodium hydroxymethylglycinate, polymethoxy bicyclic oxazolidine, dimethoxane, thimersal dichlorobenzyl alcohol, captan, chlorphenenesin, dichlorophene, chlorbutanol, glyceryl laurate, halogenated diphenyl ethers, phenolic compounds, mono- and poly-
25 alkyl and aromatic halophenols, resorcinol and its derivatives, bisphenolic compounds, benzoic esters (parabens), halogenated carbanilides, 3-trifluoromethyl-4,4'-dichlorocarbanilide, and 3,3',4-trichlorocarbanilide.
3. The cleaning concentrate of claim 2 wherein the non-cationic antimicrobial agent is a
30 mono- and poly-alkyl and aromatic halophenol selected from the group p-

chlorophenol, methyl p-chlorophenol, ethyl p-chlorophenol, n-propyl p-chlorophenol, n-butyl p-chlorophenol, n-amyl p-chlorophenol, sec-amyl p-chlorophenol, n-hexyl p-chlorophenol, cyclohexyl p-chlorophenol, n-heptyl p-chlorophenol, n-octyl p-chlorophenol, o-chlorophenol, methyl o-chlorophenol, ethyl o-chlorophenol, n-propyl o-chlorophenol, n-butyl o-chlorophenol, n-amyl o-chlorophenol, tert-amyl o-chlorophenol, n-hexyl o-chlorophenol, n-heptyl o-chlorophenol, o-benzyl p-chlorophenol, o-benzyl-m-methyl p-chlorophenol, o-benzyl-m, m-dimethyl p-chlorophenol, o-phenylethyl p-chlorophenol, o-phenylethyl-m-methyl p-chlorophenol, 3-methyl p-chlorophenol, 3,5-dimethyl p-chlorophenol, 6-ethyl-3-methyl p-chlorophenol, 6-n-propyl-3-methyl p-chlorophenol, 6-iso-propyl-3-methyl p-chlorophenol, 2-ethyl-3,5-dimethyl p-chlorophenol, 6-sec-butyl-3-methyl p-chlorophenol, 2-iso-propyl-3,5-dimethyl p-chlorophenol, 6-diethylmethyl-3-methyl p-chlorophenol, 6-iso-propyl-2-ethyl-3-methyl p-chlorophenol, 2-sec-amyl-3,5-dimethyl p-chlorophenol 2-diethylmethyl-3,5-dimethyl p-chlorophenol, 6-sec-octyl-3-methyl p-chlorophenol, p-chloro-m-cresol, p-bromophenol, methyl p-bromophenol, ethyl p-bromophenol, n-propyl p-bromophenol, n-butyl p-bromophenol, n-amyl p-bromophenol, sec-amyl p-bromophenol, n-hexyl p-bromophenol, cyclohexyl p-bromophenol, o-bromophenol, tert-amyl o-bromophenol, n-hexyl o-bromophenol, n-propyl-m,m-dimethyl o-bromophenol, 2-phenyl phenol, 4-chloro-2-methyl phenol, 4-chloro-3-methyl phenol, 4-chloro-3,5-dimethyl phenol, 2,4-dichloro-3,5-dimethylphenol, 3,4,5,6-terabromo-2-methylphenol, 5-methyl-2-pentylphenol, 4-isopropyl-3-methylphenol, para-chloro-meta-xilenol, dichloro meta xilenol, chlorothymol, and 5-chloro-2-hydroxydiphenylmethane,

4. The composition of claim 1 wherein the water soluble organic solvent is selected from C1-4 alcohols, glycol ethers, and mixtures thereof.
5. The composition of claim 1 wherein the anionic soap surfactant is selected from alkali metal soap fatty acids containing from about 8 to about 24 carbon atoms, alkali metal salts, ammonium salts, amine salts, aminoalcohol salts or the magnesium salts

of one or more of the following compounds: alkyl sulfates, alkyl ether sulfates, alkylamidoether sulfates, alkylaryl polyether sulfates, alkylaryl sulfates, alkylaryl sulfonates, monoglyceride sulfates, alkylsulfonates, alkylamide sulfonates, alkylarylsulfonates, olefinsulfonates, paraffin sulfonates, alkyl sulfosuccinates, alkyl ether sulfosuccinates, alkylamide sulfosuccinates, alkyl sulfosuccinamate, alkyl sulfoacetates, alkyl phosphates, alkyl ether phosphates, acyl sarconsinates, acyl isethionates, and N-acyl taurates.

6. The composition of claim 5 wherein the anionic soap surfactant is selected from alkali metal soap fatty acids.

7. The composition of claim 1 wherein the hydrocarbon diluent is an aromatic hydrocarbon diluent selected from Shellsolv AB, Aromatic 150, Aromatic 200 (naphthalene depleted), Aromatic 200, Aromatic 100, and HAN 857.

8. The composition of claim 1 wherein the non-cationic antimicrobial agent is present in an amount of from about 0.1 to about 5 wt%, more preferably from about 0.5 to about 5 wt%.

9. The composition of claim 1 wherein the water soluble organic solvent is present in an amount of from about 0.5 to about 10 wt%.

10. The composition of claim 1 wherein the anionic soap surfactant is present in an amount of from about 1 to about 20 wt%, more preferably from about 2 to about 15 wt%.

11. The composition of claim 1 wherein the hydrocarbon diluent is present in an amount of from about 0.1 to about 10 wt%, more preferably from about 0.5 to about 5 wt%.

12. The composition of claim 1 wherein the pine oil is present in an amount of from about 0.1 to about 15 wt%, more preferably from about 1 to about 10 wt%.

13. A hard surface cleaning concentrate composition comprising:

- a) from about 0.1 to about 5 wt% of a non-cationic antimicrobial agent;
- b) from about 0.5 to about 10 wt% of a water soluble organic solvent;
- c) from about 1 to about 20 wt% of an anionic soap surfactant;
- d) from about 0.1 to about 10 wt% of an hydrocarbon diluent;
- e) from about 0.1 to about 15 wt% of pine oil which is at least 60% terpene alcohols;
- f) optionally, from about 0 to about 10 wt% of optional materials selected from dyes, colorants, pH stabilizers and buffers, non-ionic surfactants, fragrance/fragrance enhancers, viscosity modifiers, insect repellants, and light stabilizers; and
- g) the balance being water.

14. A hard surface cleaning concentrate composition comprising:

- a) from about 0.5 to about 5 wt% of a non-cationic antimicrobial agent;
- b) from about 0.5 to about 10 wt% of a water soluble organic solvent;
- c) from about 2 to about 15 wt% of an anionic soap surfactant;
- d) from about 0.5 to about 5 wt% of an hydrocarbon diluent;
- e) from about 1 to about 10 wt% of pine oil which is at least 60% terpene alcohols;
- f) optionally, from about 0 to about 10 wt% of optional materials selected from dyes, colorants, pH stabilizers and buffers, non-ionic surfactants, fragrance/fragrance enhancers, viscosity modifiers, insect repellants, and light stabilizers; and
- g) the balance being water.

15. The cleaning concentrate of claims 1 to 14 which contain one or more insect repellants.

16. The cleaning concentrate of claim 15 wherein the amount of one or more insect repellants ranges from about 0.1 to about 5 wt%.
17. A hard surface cleaning concentrate composition comprising:
- a) from about 0.05 to about 10 wt% of a non-cationic antimicrobial agent;
 - b) from about 1 to about 20 wt% of a water soluble organic solvent;
 - c) from about 1 to about 20 wt% of an anionic soap surfactant;
 - d) from about 1 to about 15 wt% of a hydrocarbon diluent;
 - e) from about 0.001 to about 20 wt% of pine oil which is at least 60% terpene alcohols;
 - f) from about 0.1 to about 5 wt% of one or more insect repellants; and
 - g) the balance being water.
18. The cleaning concentrate of claim 17 wherein the non-cationic antimicrobial agent is selected from pyrithiones, dimethyldimethylol hydantoin, methylchloroisothiazolinone/methylisothiazolinone sodium sulfite, sodium bisulfite, imidazolidinyl urea, diazolidinyl urea, benzyl alcohol, 2-bromo-2-nitropropane-1,3-diol, formalin (formaldehyde), iodopropenyl butylcarbamate, chloroacetamide, methanamine, methyldibromonitrile glutaronitrile, glutaraldehyde, 5-bromo-5-nitro-1,3-dioxane, phenethyl alcohol, o-phenylphenol/sodium o-phenylphenol, sodium hydroxymethylglycinate, polymethoxy bicyclic oxazolidine, dimethoxane, thimersal dichlorobenzyl alcohol, captan, chlorphenenesin, dichlorophene, chlorbutanol, glyceryl laurate, halogenated diphenyl ethers, phenolic compounds, mono- and poly-alkyl and aromatic halophenols, resorcinol and its derivatives, bisphenolic compounds, benzoic esters (parabens), halogenated carbanilides, 3-trifluoromethyl-4,4'-dichlorocarbanilide, and 3,3',4-trichlorocarbanilide.
19. The cleaning concentrate of claim 18 wherein the non-cationic antimicrobial agent is a mono- and poly-alkyl and aromatic halophenol selected from the group p-chlorophenol, methyl p-chlorophenol, ethyl p-chlorophenol, n-propyl p-chlorophenol,

n-butyl p-chlorophenol, n-amyl p-chlorophenol, sec-amyl p-chlorophenol, n-hexyl p-chlorophenol, cyclohexyl p-chlorophenol, n-heptyl p-chlorophenol, n-octyl p-chlorophenol, o-chlorophenol, methyl o-chlorophenol, ethyl o-chlorophenol, n-propyl o-chlorophenol, n-butyl o-chlorophenol, n-amyl o-chlorophenol, tert-amyl o-chlorophenol, n-hexyl o-chlorophenol, n-heptyl o-chlorophenol, o-benzyl p-chlorophenol, o-benzyl-m-methyl p-chlorophenol, o-benzyl-m, m-dimethyl p-chlorophenol, o-phenylethyl p-chlorophenol, o-phenylethyl-m-methyl p-chlorophenol, 3-methyl p-chlorophenol, 3,5-dimethyl p-chlorophenol, 6-ethyl-3-methyl p-chlorophenol, 6-n-propyl-3-methyl p-chlorophenol, 6-iso-propyl-3-methyl p-chlorophenol, 2-ethyl-3,5-dimethyl p-chlorophenol, 6-sec-butyl-3-methyl p-chlorophenol, 2-iso-propyl-3,5-dimethyl p-chlorophenol, 6-diethylmethyl-3-methyl p-chlorophenol, 6-iso-propyl-2-ethyl-3-methyl p-chlorophenol, 2-sec-amyl-3,5-dimethyl p-chlorophenol 2-diethylmethyl-3,5-dimethyl p-chlorophenol, 6-sec-octyl-3-methyl p-chlorophenol, p-chloro-m-cresol, p-bromophenol, methyl p-bromophenol, ethyl p-bromophenol, n-propyl p-bromophenol, n-butyl p-bromophenol, n-amyl p-bromophenol, sec-amyl p-bromophenol, n-hexyl p-bromophenol, cyclohexyl p-bromophenol, o-bromophenol, tert-amyl o-bromophenol, n-hexyl o-bromophenol, n-propyl-m,m-dimethyl o-bromophenol, 2-phenyl phenol, 4-chloro-2-methyl phenol, 4-chloro-3-methyl phenol, 4-chloro-3,5-dimethyl phenol, 2,4-dichloro-3,5-dimethylphenol, 3,4,5,6-terabromo-2-methylphenol, 5-methyl-2-pentylphenol, 4-isopropyl-3-methylphenol, para-chloro-meta-xyleneol, dichloro meta xyleneol, chlorothymol, and 5-chloro-2-hydroxydiphenylmethane,

20. The composition of claim 1 wherein the water soluble organic solvent is selected from C1-4 alcohols, glycol ethers, and mixtures thereof.

21. The composition of claim 17 wherein the anionic soap surfactant is selected from alkali metal soap fatty acids containing from about 8 to about 24 carbon atoms, alkali metal salts, ammonium salts, amine salts, aminoalcohol salts or the magnesium salts of one or more of the following compounds: alkyl sulfates, alkyl ether sulfates,

alkylamidoether sulfates, alkylaryl polyether sulfates, alkylaryl sulfates, alkylaryl sulfonates, monoglyceride sulfates, alkylsulfonates, alkylamide sulfonates, alkylarylsulfonates, olefinsulfonates, paraffin sulfonates, alkyl sulfosuccinates, alkyl ether sulfosuccinates, alkylamide sulfosuccinates, alkyl sulfosuccinamate, alkyl sulfoacetates, alkyl phosphates, alkyl ether phosphates, acyl sarconsinates, acyl isethionates, and N-acyl taurates.

22. The composition of claim 21 wherein the anionic soap surfactant is selected from alkali metal soap fatty acids.

23. The composition of claim 17 wherein the hydrocarbon diluent is an aromatic hydrocarbon diluent selected from Shellsolv AB, Aromatic 150, Aromatic 200 (naphthalene depleted), Aromatic 200, Aromatic 100, and HAN 857.

24. The composition of claim 17 wherein the non-cationic antimicrobial agent is present in an amount of from about 0.1 to about 5 wt%, more preferably from about 0.5 to about 5 wt%.

25. The composition of claim 17 wherein the water soluble organic solvent is present in an amount of from about 0.5 to about 10 wt%.


26. The composition of claim 17 wherein the anionic soap surfactant is present in an amount of from about 1 to about 20 wt%, more preferably from about 2 to about 15 wt%.

27. The composition of claim 17 wherein the hydrocarbon diluent is present in an amount of from about 0.1 to about 10 wt%, more preferably from about 0.5 to about 5 wt%.

28. The composition of claim 17 wherein the pine oil is present in an amount of from about 0.1 to about 15 wt%, more preferably from about 1 to about 10 wt%.

29. A hard surface cleaning concentrate composition comprising:
- a) from about 0.1 to about 5 wt% of a non-cationic antimicrobial agent;
 - b) from about 0.5 to about 10 wt% of a water soluble organic solvent;
 - 5 c) from about 1 to about 20 wt% of an anionic soap surfactant;
 - d) from about 0.1 to about 10 wt% of an hydrocarbon diluent;
 - e) from about 0.1 to about 15 wt% of pine oil which is at least 60% terpene alcohols;
 - f) from about 0.1 to about 5 wt% of one or more insect repellants; and
 - 10 g) the balance being water.
30. A hard surface cleaning concentrate composition comprising:
- a) from about 0.5 to about 5 wt% of a non-cationic antimicrobial agent;
 - b) from about 0.5 to about 10 wt% of a water soluble organic solvent;
 - 15 c) from about 2 to about 15 wt% of an anionic soap surfactant;
 - d) from about 0.5 to about 5 wt% of an hydrocarbon diluent;
 - e) from about 1 to about 10 wt% of pine oil which is at least 60% terpene alcohols;
 - f) from about 0.1 to about 5 wt% of one or more insect repellants; and
 - 20 g) the balance being water.
31. The cleaning concentrate of claims 17 to 30 wherein the one or more insect repellents is selected from essential oils selected from oils of anise, citrus, aniseed, roses, mint, camphor, lemon, orange, rosemary, wintergreen, thyme, lavender, cloves, hops, tea
- 25 tree, citronella, wheat, barley, lemongrass, cedar leaf, cedarwood, cinnamon, fleagrass, geranium, sandalwood, violet, cranberry, eucalyptus, vervain, peppermint, gum benzoin, basil, fennel, fir, balsam, menthol, ocmea origanum, hydastis carradensis, berberidaceae daceae, ratanhiae, curcuma longa, Mentha arvensis (Commint), Mentha spicata (American Spearmint), Mentha cardica (Scotch
- 30 Spearmint), (-)-Limonene, (+)-Limonene, (-)-Carvone, Linalool, Alpha and Beta -

Terpineol, Fencholic acid, Borneol iso Borneol, Bornyl acetate and iso Bornyl acetate and related chemical components of the plant oils selected from anethol, catechole, camphene, pinocarvone, cedrol, thymol, eugenol, eucalyptol, ferulic acid, farnesol, hinokitiol, tropolone, limonene, menthol, methyl salicylate, carvacol, terpeneol, verbenone, berberine, ratanhia extract, caryophellene oxide, citronellic acid, curcumin, nerolidol and geraniol; N,N-diethyl-m-toluamide, diethyl phthalate, dimethyl phthalate, dibutyl phthalate, 2-Hydroxyethyl-n-octyl sulfide, N-Octyl bicycloheptene dicarboximide, Hexahydrodibenzofuran carboxaldehyde, Di-n-propyl isocinchomerate, 2-Ethyl-1,3-hexanediol, 2-(n-butyl)-2-ethyl-1,3-propanediol, Dibutyl succinate, Piperonyl butoxide, Pyrethrum, fragrances, n-alkylneoalkanamides having a formula of



wherein R₁, R₂, and R₃ are alkyl groups and the sum of the carbon atoms therein is from 6 to 12, and wherein R₄ is either a hydrogen atom or an alkyl group having one or two carbon atoms and wherein R₅ is an alkyl group having one to three carbon atoms.

INTERNATIONAL SEARCH REPORT

Int l Application No

P B 03/04422

A. CLASSIFICATION OF SUBJECT MATTER

IPC 7 C11D9/26 C11D9/24 C11D10/04 C11D3/18 C11D3/20

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 7 C11D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

EPO-Internal, WPI Data, PAJ

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	GB 2 304 115 A (RECKITT & COLMAN INC) 12 March 1997 (1997-03-12) page 1, line 1-18 page 3, line 3-14 page 9, line 3 -page 11, line 11 page 14, line 19-22 page 19, line 6-11; claims 1,3,8 ---	1-31
X	WO 97 06230 A (RECKITT & COLMAN INC) 20 February 1997 (1997-02-20) page 1, line 6,7 page 2, line 29 page 13, line 9-15 page 9, line 27 -page 10, line 16 page 15, line 23-34; tables 1B,1C --- -/-	1-31

☒ Further documents are listed in the continuation of box C.☒ Patent family members are listed in annex.

* Special categories of cited documents:

- *A* document defining the general state of the art which is not considered to be of particular relevance
- *E* earlier document but published on or after the international filing date
- *L* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- *O* document referring to an oral disclosure, use, exhibition or other means
- *P* document published prior to the international filing date but later than the priority date claimed

T later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

X document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

Y document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.

& document member of the same patent family

Date of the actual completion of the international search

16 January 2004

Date of mailing of the international search report

04/02/2004

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,
Fax: (+31-70) 340-3016

Authorized officer

Pentek, E BEST AVAILABLE COPY

INTERNATIONAL SEARCH REPORT

Int. l. Application No

P. GB 03/04422

C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 4 589 994 A (MOSEMAN ROGER E) 20 May 1986 (1986-05-20) column 1, line 59 -column 3, line 21; claims 1,6-8; examples -----	1-31
A	GB 1 236 594 A (ASPRO-NICHOLAS LIMITED) 23 June 1971 (1971-06-23) claims 1,4; examples 1,2 -----	1-31
A	US 5 763 468 A (LAUILHE JEAN-PAUL ET AL) 9 June 1998 (1998-06-09) column 3, line 44-50; claims 1,5; examples 1B,2 -----	1-31
BEST AVAILABLE COPY		

INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

P B 03/04422

Patent document cited in search report		Publication date	Patent family member(s)	Publication date
GB 2304115	A	12-03-1997	GB 2304111 A	12-03-1997
			AU 700516 B2	07-01-1999
			AU 6687096 A	05-03-1997
			BR 9609984 A	12-01-1999
			CA 2228427 A1	20-02-1997
			CN 1346878 A	01-05-2002
			CN 1198772 A ,B	11-11-1998
			DE 69628302 D1	26-06-2003
			EP 0842248 A1	20-05-1998
			ES 2195003 T3	01-12-2003
			NZ 315213 A	29-11-1999
			WO 9706230 A1	20-02-1997
			US 5591708 A	07-01-1997
			US 5728672 A	17-03-1998
			ZA 9606584 A	27-02-1997
WO 9706230	A	20-02-1997	GB 2304111 A	12-03-1997
			GB 2304112 A	12-03-1997
			AU 700516 B2	07-01-1999
			AU 6687096 A	05-03-1997
			BR 9609984 A	12-01-1999
			CA 2228427 A1	20-02-1997
			CN 1346878 A	01-05-2002
			CN 1198772 A ,B	11-11-1998
			DE 69628302 D1	26-06-2003
			EP 0842248 A1	20-05-1998
			ES 2195003 T3	01-12-2003
			GB 2304115 A ,B	12-03-1997
			NZ 315213 A	29-11-1999
			WO 9706230 A1	20-02-1997
			US 5591708 A	07-01-1997
			US 5728672 A	17-03-1998
			ZA 9606584 A	27-02-1997
			GB 2304728 A ,B	26-03-1997
			US 5629280 A	13-05-1997
			US 5939374 A	17-08-1999
US 4589994	A	20-05-1986	US 4668419 A	26-05-1987
GB 1236594	A	23-06-1971	IE 33112 B1	20-03-1974
US 5763468	A	09-06-1998	FR 2727289 A1	31-05-1996
			AT 190197 T	15-03-2000
			AU 4265196 A	19-06-1996
			CA 2181940 A1	06-06-1996
			DE 69515468 D1	13-04-2000
			DE 69515468 T2	26-10-2000
			EP 0748162 A1	18-12-1996
			ES 2145313 T3	01-07-2000
			WO 9616548 A1	06-06-1996

BEST AVAILABLE COPY